Rhizomarasmius pyrrhocephalus (Berkeley) R.H. Petersen











OVERALL: Gorman Stream Valley NRA, 10/2/2016, *J. Solem.* TOP: Gorman Stream Valley NRA, 10/2/2016, *J. Solem.* OVERALL (gills and stalk): Gorman Stream Valley NRA, 10/2/2016, *J. Solem.* GILLS and STALK: Gorman Stream Valley NRA, 10/2/2016, *J. Solem.* 

SIDE: Schooley Mill Park, 10/17/2013, J. Solem

Fire Cap Marasmius

Rhizomarasmius pyrrhocephalus Marasmiaceae

J F M A M J J A S O N D

**ID**: Cap orangish-yellow; convex, translucent. Stalk white at apex, darkens to base.

**Habitat**: Scattered, groups. Decaying oak leaves and humus under hardwoods.

**Cap**: 1/8 - 1 " [4 – 24 mm]

Orangish-yellow/brownish-orange. Moist or dry, translucent striate. Convex/rounded and smooth (broadly convex and weakly grooved in age; center often depressed).

**Gills**: White (dull pale yellow in age). Attached/notched. Close/subdistant. Many secondary gills.

Spores: White. Oval (one end slightly pointed), smooth, hyaline.

**Stalk**: 1.4 - 4" x 1/32 - 3/32" [3.5 - 10 cm x 0.5 - 2 mm] White/pale yellow at apex, progressively darkening to blackish-brown at base. Dry. Tough, Velvety. Matted fibrils on lower section. Often deeply rooted.

Frequency: Occasional.

Locations: GONRA, MURHL, SLMLP.

**Notes**: Mycobank 467533. Gorman Stream Valley NRA specimen identified by examination of spores by R. Solem. Cap of one fruiting body was very light (upper left photo).

Synonym: Marasmius pyrrhocephalus

References: BBF 201, 300. BBH 131fn. K&M 335. Kuo. Roo 184.



SPORES (7.3-8.3 x 3.6-4.1  $\mu$ m): Gorman Stream Valley NRA,10/2/2016, *R. Solem*.