

None

Russula pectinatoides Peck



OVERALL: Long Corner, PRSP, 6/27/2009, *J. Solem*.
 TOP: Franciscan Friars, 9/5/2014, *J. Solem*.
 TOP (w/ bumps on margin): Big Branch, 8/4/2014, *J. Solem*.
 OVERALL: Wincopin Trail, 8/21/2011, *J. Solem*.
 GILLS: Big Branch, 8/4/2014, *J. Solem*.
 SPORES (7.0 – 8.1 X 5.8 – 6.5 µm): Big Branch, 8/4/2014, *R. Solem*.

None

[*Russula pectinatoides*](#)
Russulaceae

J	F	M	A	M	J	J	A	S	O	N	D
					++	+++	+++	+++			

ID: Yellowish-brown, viscid, striate margin w/ bumps.

Habitat: Solitary, scattered, groups. On ground in mixed forests.

Cap: 1.2 – 3.1" [3 – 8 cm]

Yellowish-brown/straw-colored/cinnamon-brown. Convex/centrally-depressed. Margin striate w/ bumps. Viscid (wet). Skin peels.

Gills: White/pale cream. Thin. Gills taste acid.

Spores: *Cream*. Oval w/ spines 0.4-0.6 µm.

Stalk: 1 – 2" x 0.2 – 0.4" [2.5 – 5 cm x 5 – 10 mm]

White/pale yellowish (brown when bruised). Flesh white. Odor oily; taste acid.

Frequency: Very common.

Locations: BIGBR, BRNBR, CEDLP, CASCT, DANPV, FRANF, GONRA, GUILP, HANVL, MANWD, MPENA, ROGOR, SLMLP, TUFTS, UMDCF, WSTRP, WINTR.

Notes: Mycobank 213893. Big Branch and Franciscan Friars specimens identified by examination of spores by R. Solem.

References: Bni 51. BRBD 191fn. K&M 345. Kuo. Phi 147. Rog.