

Late Fall Oyster

Sarcomyxa serotina
(Persoon) V. Papp



OVERALL: Long Corner, PRSP, 11/9/2009, J. Solem.
 SIDE w/ GILLS: Long Corner, PRSP, 11/9/2009, J. Solem.
 GILLS and STALKS: Long Corner, PRSP, 11/9/2009, J. Solem.
 OVERALL: Long Corner, PRSP, 11/9/2009, R. Solem

Late Fall Oyster

[*Sarcomyxa serotina*](#)
Mycenaceae

J	F	M	A	M	J	J	A	S	O	N	D
-+								+-	---	+++	+-

ID: Cap greenish-yellow w/ purple tint, slippery/sticky (fuzzy in age). Stalk hairy, short, lateral.

Habitat: Groups, overlapping clusters. On dead hardwood logs or sticks (beech), occasionally conifers (hemlock).

Cap: 1 – 4" [2.5 – 10 cm]
 Yellowish-green/greenish-yellow, usually w/ purplish tint. Smooth/slippy/sticky (velvety/fuzzy in age). Hemispheric. Margin inrolled at first. Flesh white, firm. Taste may be bitter.

Gills: Yellowish/orangish-yellow. Attached/decurrent. Close.

Spores: Yellowish.

Stalk: Lateral. Yellowish/brownish. Hairy. Short.

Frequency: Fairly common..

Locations: BRNBR, DANPV, HENPV, HTWMA, LONGC, RKBRP.

Notes: Mycobank 433470. After autumn's first frosts.

Synonym: *Panellus serotinus* (Persoon) Kuhner

References: Bar 293. BBF 218, 304. Bin 381. Bni 215. E&S 239. Lin 789, 498. M&M 140. Phi 245. Stu 120. Roo 22.