

Bleeding Oak Crust

Stereum gausapatum (Fries) Fries



OVERALL: MPENA, 8/19/2010, R. Orr.

FERTILE SURFACE (with red drops): MPENA, 8/19/2010, R. Orr.

INFERTILE SURFACE: MPENA, 8/19/2010, J. Solem.

END OF LOG w/ FRUITING BODIES: MPENA, 8/19/2010, J. Solem.

Bleeding Oak Crust

Stereum gausapatum

Stereaceae

J	F	M	A	M	J	J	A	S	O	N	D	
-----							++	-----				

ID: Zonate bands of orange/brown/black; wavy margin.

Habitat: Dead or fallen oaks.

Fruiting Body: Up to 1.5" x 1.5" [4 cm x 4 cm]

Bands of orange/brown/black, zonate. Edges usually curl up. Margin very light when fresh. Flexible (brittle when dry). Often w/ sweet almond odor.

Fruiting Surface: Orange shades tinged w/ brown. Smooth/wrinkled. Turns red where injured. May ooze red drops.

Spores: Pale orange.

Frequency: Uncommon.

Locations: MPENA, WSTRP.

Notes: Mycobank 193672.

References: BBF 439key. Bin 419. Rog. W&L 489fn.

https://www.messiah.edu/Oakes/fungi_on_wood/crust%20and%20parchment/species%20pages/Stereum%20gausapatum.htm