
Hard's Stropharia

Stropharia hardii G.F. Atkinson



GILLS, STALK and CAP: MPENA,10/13/2010, J. Solem. GILLS, RING, STALKS and RHIZOMORPHS: MPENA,10/13/2010, J. Solem.

Hard's Stropharia



ID: Ochre cap, darker in center. Persistent ring.

Habitat: Single, scattered. On woody debris in hardwood forests; also wood chips.

Cap: 1 – 4" [2.5 – 10 cm]

Ochre/brownish-ochre (yellowish-ochre in age, darker in center). Smooth (covered w/matted fibrils/scales in age). Convex (nearly flat in age). Flesh whitish.

Gills: Whitish (purple-brown in age). Attached. Close.

Spores: *Purple-brown*. Ellipsoid, smooth w/ apical pore.

Stalk: 1.5 - 4.8" x $\frac{1}{4} - \frac{3}{4}$ " [4 - 12 cm x 6 - 20 mm] Whitish/ochre-yellow. Slightly scaly/nearly smooth. Sticky (dry in age). Nearly equal. Persistent superior ring. Base w/ whitish rhizomorphs.

Frequency: Uncommon.

Locations: MPENA, PIGTL, WSKPV.

Notes: Mycobank 217638. MPENA record identified by examination of spores by R. Solem.

References: BBF 250, 310. BBH188. Bni 333. K&M 363. K&M 363. Kuo. Lin 728, 202. M&M 255fn. Phi 227. Roo 77. Stu 218.



SPORES (5.4-6.2 X 3.6-4.3 µm): MPENA, 10/13/2010, R. Solem.