

Slippery Jack
Sticky Bun

Suillus luteus
(Linnaeus) Roussel



CAP: Schooley Mill Park, 11/9/2017, *J. Solem*.
GILLS, STALK, and RING: Schooley Mill Park, 11/9/2017, *J. Solem*.
CAP (variant): Schooley Mill Park, 11/9/2017, *J. Solem*.
STALK, PARTIAL VEIL, CAP: Schooley Mill Park, 11/9/2017, *J. Solem*.
SPLIT: Schooley Mill Park, 11/9/2017, *J. Solem*.

Slippery Jack
Sticky Bun

Suillus luteus
Boletaceae

J	F	M	A	M	J	J	A	S	O	N	D
						---	---	---	---	++	

ID: Some shade of brown w/ glutinous cap. Prominent ring.

Habitat: Solitary, groups. Under pines.

Cap: 2 – 4.7" [5 – 12 cm]

Reddish-/cinnamon-/yellow-brown or ocher. Convex, often w/ low umbo (broadly convex/flat in age). Glutinous/viscid when fresh, sticky/shiny when dry. Margin w/ sterile tissue, often w/ remnants of partial veil. Flesh white/pale yellow; thick in center.

Pores: Whitish/pale yellow (yellow to dark-/olive-yellow in age)

Depressed at stalk in age. Angular (1-2/mm). Tubes (4-15 mm). When young, covered w/ white partial veil w/ dark-/grayish-purple tones.

Spores: *Orangish-brown*. Fusiform, smooth.

Stalk: 1.1 – 3.1" X 0.4 – 1" [3 – 8 cm X 1 – 2.5 cm]

White (pale yellow in age at apex, often w/ purplish/brownish tones toward base). Resinous dots/smears above ring. Flaring/sleeve-like ring w/ purple tones on underside.

Frequency: Uncommon.

Locations: SLMLP.

Notes: Mycobank 120823. Schooley Mill Park specimen identified by examination of spores by R. Solem

References: Bar 170. BBF 356, 370. BBH 215. Bni 407.

BRB1 243, 337. BRB2 341-3. K&M 367. Kuo. Lin 586, 401. M&M 364. M&S 176. Phi 288. Roo 287. Stu 294.



SPORES (7.4-8.0 x 3.0-3.6 µm): Schooley Mill Park, 11/9/2017, *R. Solem*.