

Slippery Jill

Suillus salmonicolor (Frost) Halling



OVERALL: Rockburn Branch Park, 10/18/2010, J. Solem.

TOP: Rockburn Branch Park, 10/18/2010, J. Solem.

BUTTON STAGE and SPLIT FRUITING BODY: Rockburn Branch Park, 10/18/2010, J. Solem.

Slippery Jill

Suillus salmonicolor
Boletaceae

J	F	M	A	M	J	J	A	S	O	N	D
							---	---	---	+	---

ID: Cap varies yellow to brown, very slimy when wet; convex/nearly plane. Pores dingy yellow (brownish in age). Stalk yellow w/ reddish glandular dots. Use NH₄OH and KOH.

Habitat: Scattered, groups. Under pines; often sandy soil in moss.

Cap: 1.1 – 3.8" [3 – 9.5 cm]

Variable (dingy yellow, yellowish-orange, ochraceous-salmon, cinnamon-/olive-/yellow-brown. Dry: shiny. Wet: sticky/slimy. Bluntly rounded/convex/nearly plane. Flesh yellow/orange/orange-buff.

Pores: Yellow/dingy yellow/yellowish-orange/salmon (brownish in age). Circular/angular (1 – 2/mm). Tubes (8 – 10 mm).

Spores: Brown/cinnamon-brown.

Stalk: 1 – 4" x 1/4 – 5/8" [2.5 1/4 10 cm x 6 – 16 mm]

Whitish/yellowish/pinkish-ochraceous, speckled w/ reddish-/dark-brown glandular dots/smears. Flesh yellow to salmon-orange. Veil leaves a flaring, rubbery ring w/ white roll of tissue and lavender edge.

Frequency: Uncommon.

Locations: RKBRP.

Notes: Button and very young w/ thick sticky partial veil. Cap purplish-red, then purplish-black w/ NH₄OH or KOH. Flesh dark purplish-red w/ NH₄OH or KOH.

Synonyms: *Suillus subluteus*. *Suillus pinorigidus*. *Suillus cothurnatus*.

References Bar 171. BBF 358, 370. BBH 213fn. Bni 411.

BRB1 250, 340. BRB2 348. BRBD 238, 56. K&M 369. Lin 589, 371. M&M 365. Roo 286fn.