

None

Trametes hirsuta (Wulfen) Lloyd



OVERALL: Pigtail, 4/2/2010, R. Orr.

None

Trametes hirsuta
Polyporaceae

J	F	M	A	M	J	J	A	S	O	N	D
-	+	-	-	-	+	+	-	-	-	-	+

ID: Cap white/brown concentrically zoned; persists; zoned w/ gray-white/yellow-pale brown. Flesh thick, two-layered.

Habitat: Clusters on dead hardwoods (rarely conifers).

Fruiting Body: 1 – 4" [2.5 cm x 10 cm]

Zones w/ gray/whitish/brownish shades, little contrast. Densely hairy, radially furrowed in concentric zones. Semicircular/irregularly bracket-/kidney-shaped. Margin often brownish/blackish. Flesh ($\leq 7/16$ " [18mm]), upper layer grayish (soft, fibrous); separated from lower layer (white, tough) by thin, black line.

Pores: Whitish (brownish/grayish/yellowish in age). Pores circular/angular (3 – 4/mm). Tubes (6 mm).

Frequency: Fairly common.

Locations: BRNBR, DANPV, HIRIP, PATBT, PIGTL, ROBNC, SLMLP, WINTR.

Notes: Mycobank 267192.

Synonym: *Coriolus hirsutus*.

References: Bar 137. BBF 394fn. BBH 22. Bin 429. Bni 433fn. BRBD 271, 65. Kuo. Lin 489fn. Roo 386fn.