None

Trametopsis cervina (Schweinitz) Tomsovsky



INFERTILE SURFACE: Big Branch, 5/29/2009, *R. Solem.* PORES: Big Branch, 5/29/2009, *R. Solem.* 

None

	_							~	•		_
J	F	M	Α	M	J	J	Α	S	0	N	D
	-+-		+-+	+	++-						

**ID**: Body zones white/pinkish-buff; s-/fan-shaped, overlap- ping clusters; coarse hairs, margin uneven/wavy in age; stalk- less. Pores white (buff in age), labyrinthine/tooth-like in age.

Habitat: Overlapping clusters on decaying hardwoods (rarely conifers).

**Fruiting Body**: 1.1 - 3.5" x 0.4" thick [3 - 9 cm x 9 mm]White/pinkish-buff/pale cinnamon zones. Coarse hairs. Shell-/fanshaped, leathery to stiff. Margin uneven/wavy in age. Flesh (1 mm) white/pale buff, fibrous-tough.

**Pores**: White (cinnamon-buff in age). Irregular/angular (conspicuously labyrinthine/tooth-like in age).

Spores: White/cream.

Frequency: Occasional.

Locations: BIGBR, CASCT, DANPV, PIGTL, RKBRP.

Notes: Mycobank 511830.

Synonym: Trametes cervina.

References: BBF 380, 399.