

Bay Cap Bolete

*Tylophilus badiceps* (Peck)  
A.H. Smith & Thiers



TOPS: Wilde Lake, 7/7/2015, R. Orr.  
PORES, CAPS: Wilde Lake, 7/7/2015, R. Orr.  
HABITAT: Wilde Lake, 7/7/2015, R. Orr.

Bay Cap Bolete

*Tylophilus badiceps*  
Boletaceae

J	F	M	A	M	J	J	A	S	O	N	D
						+	-	-	-	-	

**ID:** Maroon cap, white pores, stalk color of cap.

**Habitat:** Solitary, groups. On ground under hardwoods (especially oaks).

**Cap:** 1.8 – 3.5" [4.5 – 9 cm]  
Maroon to pinkish/dull reddish-brown. Convex (broadly convex/flat in age). Velvety (smoother in age). Dry. Flesh white

**Pores:** White (dingy white/brownish in age; bruises brown. Pores 1-3/mm.

**Spores:** Pinkish/deep flesh color. Narrowly oval, smooth.

**Stalk:** 2.5 – 3.5" x 0.6 – 1.1" [5 – 9 cm x 1.5 – 2.8 cm]  
Color of cap w/ white apex. Surface smooth, glabrous or w/ reticulation near apex. Equal or w/ mid-portion/base swollen. Solid.

**Frequency:** Uncommon.

**Locations:** WILLK.

**Notes:** Mycobank 325149. Wilde Lake specimen identified by examination of spores by R. Solem.

**References:** BBF 381key. BRB1 260, 341. BRB2 371. Bin 229. Phi 284.



SPORES (8.8-9.7 X 3.0-3.3 µm): Wilde Lake, 7/7/2015, R. Orr.