

Burnt-orange Bolete

*Tylophilus balloui* (Peck) Singer



OVERALL: Hammond Village, 8/8/2017, J. Solem.  
 OVERALL (old): Rocky Gorge, 9/1/2011, J. Solem.  
 PORES and STALK: Rockburn Branch, 7/28/2010, J. Solem.  
 CUT (shows bruising): Franciscan Friars, 8/2/2014, J. Solem.

Burnt-orange Bolete

*Tylophilus balloui*  
 Boletaceae

J	F	M	A	M	J	J	A	S	O	N	D
						+++	+++	+++	--		

**ID:** Bright to dull orange-red, convex (irregular in age). Pores white to tan.

**Habitat:** Solitary, scattered, groups. On ground in hardwood forests or under hardwood trees in lawns.

**Cap:** 2 – 4.8" [5 – 12 cm]  
 Bright orange/orange-red (dull orange/cinnamon/tan in age). Convex (nearly flat or irregular in age). Dry. Flesh white (bruises pinkish-tan/violet-brown).

**Pores:** White/dingy white (tan/pinkish in age). Bruise brown. Angular (1 – 2/mm).

**Spores:** Yellow/dark cream. Ellipsoid, smooth

**Stalk:** 1 – 4.8" x 1/4 – 1" [2.5 – 12 cm x 6 – 25 mm]  
 Whitish, tinged yellow/orange (bruises brownish). Smooth/scurfy. Equal or base swollen.

**Frequency:** Fairly common.

**Locations:** FRANF, GATWY, GONRA, HAMVL, MANWD, MCHPV, OTHER, ROBNC, ROGOR, RKBRP, WINTR.

**Notes:** Mycobank 516813. Franciscan Friars specimen identified by examination of spores by R. Solem.

**Synonym:** *Rubinoboletus balloui*

**References:** BBF 359, 371. BBH 220. Bni 396. BRB1 261, 342. BRB2 374. BRBD 240, 56. E&S 270. K&M 382. Kuo. Lin 591, 397. M&M 378. Phi 284. Stu 262.



SPORES (6.7 – 8.6 X 4.0 – 4.3 µm): Franciscan Friars, 8/2/2014, R. Solem..