Uncommon 48-62 mm (1.9-2.4 in.) Large MALE: Face: black. Body: dark pale-gray. Wings: black streak basal, spot in middle, bar/spot on front margin of outer half. HW: white spot basal. Legs: black. Appendages: blackish.

Bar-winged Skimmer

- FEMALE: Eyes: red-brown. Abdomen brown or lightly pruinose; S8 expanded laterally, usually brownish. Wings: some lack black basal streaks (only a tiny spot may be present mid-wing); some have smoky-brown color on tips.
- JUVENILE: Similar to L. vibrans (Great Blue S.) Face: brownish. Thorax: yellow w/ brown triangular marking on the side basal of FW; lateral pale pattern resembles a cartoon wolf. Abdomen: middorsal/lateral brown stripes w/ yellow between.

Habitat: Temporary pools, marshy edges of ponds. More often in forested habitats than other Libellula.

Reproduction: Mate perched on weed or bush. Female oviposits alone while male guards.

Similar Species: Female L. axilena, L. incesta (Slaty S.), L. vibrans (Great Blue S.)

Locations: Big Branch, Coca Cola Drive, David Force, Dunteachin Farm, Elkhorn, Farm Ponds, Font Hill, Guilford - GV, Hipsley Mill, Kindler - P, Long Corner - P, Milk Plant, Mission, MPEA - P4, Pigtail, Stevens - E/W.

MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC

Libellula axilena Westwood

30 May - 23 Aug