

White Oysterette

Cheimonophyllum haedinum
(Berkeley & M.A. Curtis)
Valade & P.-A. Moreau



OVERALL: Murray Hill,
10/30/2019, *J. Solem*.
TOP: Daniels, PVSP, 9/6/2012,
R. Solem.
GILLS: MPENA, 8/30/2012, *J. Solem*.
GILLS (mature): Daniels, PVSP,
9/6/2012, *R. Solem*.

White Oysterette

Cheimonophyllum haedinum
Cyphellaceae

J	F	M	A	M	J	J	A	S	O	N	D
						+	+	+	-	+	+

ID: Cap kidney-shaped w/ white hairs. Stalk off-center or none.

Habitat: Solitary, groups. On dead hardwoods.

Cap: 0.4 – 0.8" [1 – 2 cm]

White hairs on yellowish background. Hemispheric/kidney-shaped. Margin inrolled, slightly grooved in age. Flesh thin, soft, whitish/yellowish.

Gills: White/yellowish-tan. Decurrent. Subdistant/fairly close. Rather broad.

Spores: *White*. Round, smooth, hyaline.

Stalk: Off-center/none.


Frequency: Fairly common.

Locations: BIGBR, DANPV, FRANF, HENPV, LONGC, MPENA, MURHL, ROGOR, SYRIV, WINTR..

Notes: Mycobank 846794.

Synonym: *C. candidissimum* . *Pleurotus candidissimus*.

References: Bar 297. BBF 75, 276. Bni 218. Lin 743, 496. M&M 138. Phi 67.



SPORES (5.4-6.4 µm): Murray Hill, 10/30/2019, *R. Solem*.