Elephant Ear Gabled False Morel

Neogyromitra brunnea (Underwood) Herter



OVERALL: Wincopin Trail, 3/25/2011, G. King. OVERALL: Wincopin Trail, 3/"6/2011, R. Solem.

OVERALL: Daniels, 4/7/2012, *B. Ott.*OVERALL: Daniels, 4/7/2012, *B. Ott.*SIDE: Wincopin Trail, 3/26/2011, *R. Solem.*CUT: Wincopin Trail, 3/26/2011, *R. Solem.*

Elephant Ear Gabled False Morel Neogyromitra brunnea
Discinaceae

J F M A M J J A S O N D ++-+-

ID: Wrinkled yellow-/reddish-brown contorted cap w/ thick stalk.

Habitat: Solitary, scattered, on ground under hardwoods.

Cap: 1.5-4.8" wide x 1.5-3.5" tall [4-12 cm x 4-9]Fertile (upper) surface yellow-/reddish-brown. Lower (sterile) surface pale yellow-brown/whitish (slightly granular and often fused to part of stalk). Irregularly folded/divided into deep wrinkled lobes;can be saddle-shaped. Moist.

Stalk: 1-4" x 1.5-3.5" thick [2.5-10 cm x 4-9 cm] Chalky/pale grayish-white. Ribbed or may appear to be joined/ribbed stalks. Branched at top. Usually wider than high; equal or wider at base. Chambered interior.

Spores: White; *brown*, powdery at maturity.

Frequency: Occasional.

Locations: DANPV, MPENA, PIGTL, SAVMT, WINTR.

Notes: Mycobank 301810

Synonyms: Gyromitra brunnea, Gyromitra fastigeata.,

References: BBB 146. BBF 474, 478. BBH 348. E&S 43FN. K&M 192.

Kuo. Lin 337, 718. M&M 510. Roo 478.